

L6 ANSWER 43 OF 50 CAPLUS COPYRIGHT 2002 ACS

AN 1992:511645 CAPLUS

DN 117:111645

TI Preparation of 2-(2-pyridyl)pyrimidines as agrochemical fungicides

IN Robson, Michael John; Worthington, Paul Anthony

PA Imperial Chemical Industries PLC, UK

SO PCT Int. Appl., 34 pp.


CODEN: PIXXD2

DT Patent

LA English


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9208714	A1	19920529	WO 1991-GB1853	19911023
	W: AU, BB, BG, BR, CA, CS, FI, HU, JP, KP, KR, LK, MC, MG, MN, MW, NO, PL, RO, SD, SU, US				
	RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FR, GA, GB, GN, GR, IT, LU, ML, MR, NL, SE, SN, TD, TG				
	AU 9188527	A1	19920611	AU 1991-88527	19911023
	EP 557291	A1	19930901	EP 1991-917011	19911023
	EP 557291	B1	19960814		
	R: CH, DE, DK, FR, GB, LI				
	ZA 9108683	A	19920729	ZA 1991-8683	19911031
	US 5208239	A	19930504	US 1991-785646	19911031
PRAI	GB 1990-24873		19901115		
	WO 1991-GB1853		19911023		
OS	MARPAT 117:111645				
AB	Title compds. [I; R1 = (substituted) Ph, cycloalkyl; R2-R7 = H, alkyl, haloalkyl, alkoxy, amino; Z = C.tplbond.C, (alkyl-substituted) vinylene, ethylene] were prepd. Thus, 2,6-dibromopyridine in THF at -78.degree. was treated successively with BuLi, ZnCl2 in Et2O, (Ph3P)4Pd, and 2-bromo-4-methylpyrimidine. The mixt. was then refluxed 4 h to give 30% 2-(6-bromo-2-pyridinyl)-4-methylpyrimidine. This was stirred with CuI, phenylacetylene, and (Ph3P)2PdCl2 in THF/Et3N at 20.degree. to give 54% title compd. II. II as a 100 ppm spray gave complete control of Erysiphe graminis on wheat and of Venturia inaequalis on apples.				
IT	142790-12-3P	142790-15-6P	142790-16-7P		
	142790-17-8P	142790-18-9P	142790-19-0P		
	142790-20-3P	142790-21-4P	142790-22-5P		
	142790-23-6P				
	RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (prepn. of, as agrochem. fungicide)				
RN	142790-12-3	CAPLUS			
CN	Pyrimidine, 4-methyl-2-[6-(phenylethynyl)-2-pyridinyl]- (9CI) (CA INDEX NAME)				




RN 142790-15-6 CAPLUS

CN Pyrimidine, 2-[6-[(4-chlorophenyl)ethynyl]-2-pyridinyl]-4-methyl- (9CI) (CA INDEX NAME)




RN 142790-16-7 CAPLUS

CN Pyrimidine, 2-[6-[(2-chlorophenyl)ethynyl]-2-pyridinyl]-4-methyl- (9CI)  
(CA INDEX NAME)




RN 142790-17-8 CAPLUS

CN Pyrimidine, 2-[6-[(3-chlorophenyl)ethynyl]-2-pyridinyl]-4-methyl- (9CI)  
(CA INDEX NAME)




RN 142790-18-9 CAPLUS

CN Pyrimidine, 4-methyl-2-[6-[(4-methylphenyl)ethynyl]-2-pyridinyl]- (9CI)  
(CA INDEX NAME)




RN 142790-19-0 CAPLUS


CN Pyrimidine, 2-[6-[(3,4-dichlorophenyl)ethynyl]-2-pyridinyl]-4-methyl- (9CI) (CA INDEX NAME)




RN 142790-20-3 CAPLUS  
 CN Pyrimidine, 2-[6-[(2,3-dichlorophenyl)ethynyl]-2-pyridinyl]-4-methyl-  
 (9CI) (CA INDEX NAME)




RN 142790-21-4 CAPLUS  
 CN Pyrimidine, 2-[6-[(cyclohexylethynyl)-2-pyridinyl]-4-methyl- (9CI) (CA  
 INDEX NAME)



RN 142790-22-5 CAPLUS  
 CN Pyrimidine, 2-[6-[(1,1'-biphenyl)-4-ylethynyl]-2-pyridinyl]-4-methyl-  
 (9CI) (CA INDEX NAME)



RN 142790-23-6 CAPLUS  
 CN Pyrimidine, 2-[6-(3,3-dimethyl-1-butynyl)-2-pyridinyl]-4-methyl- (9CI)  
 (CA INDEX NAME)






File Copy

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b>  <b>C07D 401/04, A01N 43/54</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 92/08714</b>  <b>(43) International Publication Date:</b> <b>29 May 1992 (29.05.92)</b>
<b>(21) International Application Number:</b> PCT/GB91/01853 <b>(22) International Filing Date:</b> 23 October 1991 (23.10.91)  <b>(30) Priority data:</b> 9024873.3 15 November 1990 (15.11.90) GB  <b>(71) Applicant (for all designated States except US):</b> IMPERIAL CHEMICAL INDUSTRIES PLC [GB/GB]; Imperial Chemical House, Millbank, London SW1P 3JF (GB).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> ROBSON, Michael, John [GB/GB]; 12 Greenham Wood, North Lake, Bracknell, Berkshire RG12 4WJ (GB). WORTHINGTON, Paul, Anthony [GB/GB]; 4 Oakhurst, Maidenhead Court Park, Maidenhead, Berkshire SL6 8HY (GB).		<b>(74) Agent:</b> HOUGHTON, Malcolm, John; Imperial Chemical Industries plc, Legal Dept. Patents, PO Box No 6, Bessmer Road, Welwyn Garden City, Herts AL7 1HD (GB).  <b>(81) Designated States:</b> AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL (European patent), NO, PL, RO, SD, SE (European patent), SN (OAPI patent), SU <sup>+</sup> , TD (OAPI patent), TG (OAPI patent), US.  Published <i>With international search report.</i>

**(54) Title:** PYRIDIL-PYRIMIDINE DERIVATIVES AS FUNGICIDAL COMPOUNDS**(57) Abstract**

Fungicidal compounds having formula (I), wherein R<sup>1</sup> is an optionally substituted phenyl group, C<sub>3-6</sub> cycloalkyl optionally substituted by alkyl or alkyl; R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are independently hydrogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> alkoxy or NR'R''; Z is a linking group and is -C≡C-, -CR<sup>8</sup>=CR<sup>10</sup> or -CR<sup>8</sup>R<sup>9</sup>-CR<sup>10</sup>R<sup>11</sup>.; R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R' and R'' are independently hydrogen or C<sub>1-4</sub> alkyl; provided that when R is alkyl Z is not -CR<sup>8</sup>R<sup>9</sup>-CR<sup>10</sup>R<sup>11</sup>.; and acid addition salts or metal complexes thereof; and processes for preparing them, compositions comprising them and methods of using them to combat fungi.

# + DESIGNATIONS OF "SU"

Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

## *FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LJ	Liechtenstein	SU+	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

## PYRIDIL-PYRIMIDINE DERIVATIVES AS FUNGICIDAL COMPOUNDS

The present invention relates to pyridylpyrimidine derivatives that are useful as fungicides, to processes for preparing them, to fungicidal compositions containing them, and to methods of using them to combat fungi, especially fungal infections in plants.

According to the present invention there is provided a compound having the general formula (I), wherein  $R^1$  is an optionally substituted phenyl group,  $C_{3-6}$  cycloalkyl optionally substituted by alkyl, or alkyl;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  alkoxy or  $NR'R''$ ; Z is a linking group and is  $-C\equiv C-$ ,  $-CR^8=CR^{10}-$  or  $-CR^8R^9-CR^{10}R^{11}-$ ;  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{11}$ ,  $R'$  and  $R''$  are independently hydrogen or  $C_{1-4}$  alkyl; provided that when  $R^1$  is alkyl Z is not  $-CR^8R^9-CR^{10}R^{11}-$ ; and acid addition salts or metal complexes thereof.

Alkyl groups and the alkyl moiety of the alkoxy group contain from 1 to 6, especially from 1 to 4, carbon atoms and are in the form of straight or branched chains, i.e. methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl or tert-butyl.

Halogen includes fluorine, chlorine, bromine and iodine atoms.

Haloalkyl groups contain from 1 to 4 carbon atoms and at least one halogen. They are in the form of either straight or branched chains and are, for example, fluoromethyl, chloromethyl, fluoroethyl, chloroethyl, trifluoromethyl, trichloromethyl, pentafluoromethyl or hexafluoro-iso-propyl.

Cycloalkyl contains from 3 to 6 carbon atoms and is, for example, cyclopropyl, cyclopentyl or cyclohexyl.

The acid addition salts of the compounds of the invention include salts with inorganic or organic acids, for example, hydrochloric, nitric, sulphuric, acetic, 4-toluene-sulphonic or oxalic acids.

The metal complexes of the compounds of the invention include complexes with copper, zinc or manganese cations.

Optional substituents on the phenyl group include halogen (especially chlorine, fluorine or bromine),  $C_{1-4}$  alkyl (especially methyl),  $C_{1-4}$  alkoxy (especially methoxy),  $C_{1-4}$  haloalkoxy (especially trifluoromethoxy),  $C_{2-4}$  alkenyl (especially vinyl or allyl),  $C_{2-4}$  alkynyl (especially ethynyl or propargyl),  $C_{1-4}$  haloalkyl (especially trifluoromethyl or mono- or di- $(C_{1-4})$ alkylamino (especially methylamino or dimethylamino), or phenyl, benzyl, phenoxy or benzyloxy optionally substituted with halogen

(especially chlorine or fluorine),  $C_{1-4}$  alkyl (especially methyl),  $C_{1-4}$  haloalkyl (especially trifluoromethyl),  $C_{1-4}$  alkoxy (especially methoxy) or  $C_{1-4}$  haloalkoxy (especially trifluoromethoxy).

Alkenyl groups contain from 2-4 carbon atoms and are alk-1-enyl, alk-2-enyl or alk-3-enyl, for example, vinyl, 2-prop-1-enyl, 1-prop-1-enyl, allyl, 2-but-2-enyl, 1-(2-methylprop-1-enyl), 1-but-1-enyl, 1-(1-methylprop-2-enyl), 1-(2-methylprop-2-enyl), 1-but-2-enyl, or 1-but-3-enyl.

Alkynyl groups contain from 2 to 4 carbon atoms and are, for example, ethynyl, prop-1-ynyl, propargyl or 2-but-3-ynyl.

Compounds wherein Z is  $-CR^8=CR^{10}-$  can exist as geometric isomers. The present invention covers both isomers and mixtures in any proportion of these isomers.

In one particular aspect the present invention provides a compound of general formula (I) wherein  $R^1$  is an optionally substituted phenyl group (especially optionally substituted with halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkoxy,  $C_{1-4}$  alkenyl,  $C_{2-4}$  alkynyl,  $C_{1-4}$  haloalkyl or phenyl);  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  haloalkyl or  $C_{1-4}$  alkoxy; Z is a linking group and is  $-C\equiv C-$ ,  $-CR^8=CR^{10}-$  or  $-CR^8R^9-CR^{10}R^{11}-$ ;  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  are independently hydrogen or  $C_{1-4}$  alkyl; and acid addition salts or metal complexes thereof.

In another aspect the present invention provides a compound of general formula (I) wherein  $R^1$  is an optionally substituted phenyl group,  $C_{3-6}$  cycloalkyl optionally substituted by alkyl or alkenyl;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  alkoxy or  $NR'R''$ ; Z is  $-CR^8=CR^{10}-$  or  $-C\equiv C-$ ;  $R^8$  and  $R^{10}$  are independently  $C_{1-4}$  alkyl or, preferably, hydrogen;  $R'$  and  $R''$  are independently hydrogen or  $C_{1-4}$  alkyl; and acid addition salts or metal complexes thereof.

In a further aspect the present invention provides a compound of general formula (I) wherein  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$  and  $R^7$  are all hydrogen and  $R^5$  is  $C_{1-4}$  alkyl (especially methyl).

In a still further aspect the present invention provides a compound of general formula (I) wherein  $R^1$  is phenyl optionally substituted with halogen (especially fluorine, chlorine or bromine),  $C_{1-4}$  alkyl (especially methyl),  $C_{1-4}$  alkoxy (especially methoxy),  $C_{1-4}$  haloalkoxy (especially trifluoromethoxy),  $C_{2-4}$  alkenyl (especially vinyl and allyl),  $C_{2-4}$  alkynyl (especially ethynyl and propargyl),  $C_{1-4}$  haloalkyl (especially trifluoromethyl) or phenyl; Z is  $-C\equiv C-$ ;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$  and  $R^7$  are all


hydrogen;  $R^5$  is  $C_{1-4}$  alkyl (especially methyl); and acid addition salts or metal complexes thereof.

In yet another aspect the present invention provides a compound of general formula (I) wherein  $R^1$  is phenyl optionally substituted with halogen (especially chlorine),  $C_{1-4}$  alkyl (especially methyl),  $C_{1-4}$  haloalkoxy (especially trifluoromethoxy) or phenyl, or  $R^1$  is cyclohexyl; Z is  $-CH=CH-$  or  $-C\equiv C-$ ;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$  and  $R^7$  are all hydrogen; and  $R^5$  is  $C_{1-4}$  alkyl (especially methyl).

In another aspect the invention provides compounds of formula (I) in which  $R^1$  is a phenyl group,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$  and  $R^7$  are all hydrogen,  $R^5$  is  $C_{1-4}$  alkyl (especially methyl) and Z is  $-C\equiv C-$  or  $-CH=CH-$ .

Examples of compounds of the invention of formula (I) are given in Table I.

TABLE I



Compound No	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	Z	mp °C	Comments
1	C <sub>6</sub> H <sub>5</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-	83-85	
2	C <sub>6</sub> H <sub>5</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-	oil	trans:cis 1:2
3	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-	130-131	
4	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-		
5	2,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-		
6	2-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-	114-116	
7	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-		

TABLE I (continued)

Compound No	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	Z	mp °C	Comments
8	2-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-		
9	3-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-	116-117	
10	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-	160-161	
11	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-		
12	4-CF <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-		
13	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-		
14	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
15	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
16	2,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	H	H	H	-C≡C-		
17	2-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
18	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
19	2-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
20	3-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
21	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
22	4-CF <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
23	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
24	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-C≡C-		
25	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
26	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis

TABLE I (continued)

Compound No	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	Z	mp °C	Comments
27	2,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
28	2-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
29	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
30	2-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
31	3-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
32	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
33	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
34	4-CF <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
35	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		cis
36	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
37	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
38	2,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
39	2-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
40	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
41	2-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
42	3-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
43	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
44	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
45	4-CF <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans

TABLE I (continued)

Compound No	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	Z	mp °C	Comments
46	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH=CH-		trans
47	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
48	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
49	2,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	H	H	H	-CH=CH-		cis
50	2-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
51	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
52	2-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
53	3-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
54	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
55	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
56	4-CF <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
57	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		cis
58	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
59	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
60	2,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	H	H	H	-CH=CH-		trans
61	2-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
62	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
63	2-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans

TABLE I (continued)

Compound No	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	Z	mp °C	Comments
64	3-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
65	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
66	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
67	4-CF <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
68	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH=CH-		trans
69	C <sub>6</sub> H <sub>5</sub>	H	H	H	H	H	H	-CH=CH-		cis
70	C <sub>6</sub> H <sub>5</sub>	H	H	H	H	H	H	-CH=CH-		trans
71	C <sub>6</sub> H <sub>5</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
72	C <sub>6</sub> H <sub>5</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
73	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
74	4-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
75	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
76	4-F-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
77	2,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
78	2,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
79	2-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
80	2-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
81	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		

TABLE I (continued)

Compound No	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	Z	mp °C	Comment
82	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
83	2-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
84	2-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
85	3-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
86	3-Cl-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
87	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
88	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
89	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
90	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
91	4-CF <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
92	4-CF <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
93	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	H	H	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
94	4-Br-C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -		
95	3,4-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	CH <sub>3</sub>	H	H	-CH <sub>2</sub> -CH <sub>2</sub> -	137-139	
96	2,3-Cl <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-	119-120.5	
97	cyclohexyl	H	H	H	CH <sub>3</sub>	H	H	-C≡C-	107.5-108.5	
98	4-C <sub>6</sub> H <sub>5</sub> -C <sub>6</sub> H <sub>4</sub>	H	H	H	CH <sub>3</sub>	H	H	-C≡C-	176-179°C	
99	tert-butyl	H	H	H	Me	H	H	-C≡C-	98-100°C	
100	(CH <sub>3</sub> ) <sub>3</sub> Si	H	H	H	Me	H	H	-C≡C-		

TABLE II : SELECTED PROTON NMR DATA

Table II shows selected proton NMR data for certain compounds described in Table I along with mass spectral data which is the molecular ion. Chemical shifts are measured in ppm from tetramethylsilane, and deuteriochloroform was used as a solvent throughout.

The following abbreviations are used:

dd = doublet of doublets      t = triplet  
s = singlet                      m = multiplet  
d = doublet

Compound No	NMR DATA
1	2.66(3H,s); 7.19(1H,d); 7.35(3H,m); 7.62(3H,m); 7.85(1H,t); 8.45(1H,d); 8.78(1H,d) ppm. $M^+ = 271$
2	2.68(3H,two singlets); 7.18(1H,d); 7.25-7.40(4H,m); 7.58-7.70(4H,m); 7.85(1H,t); 8.33-8.46(1H,dd); 8.80(1H,dd) ppm. $MH^+ = 274$

Compounds of formula (I), wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are as defined above and Z is an acetylene ( $-C\equiv C-$ ) linkage, can be prepared by treating a halopyridine of general formula (II), wherein  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are as defined above and Hal is a halogen (preferably bromine or iodine), with an acetylene of general formula (III), wherein  $R^1$  is as defined above, in the presence of a convenient palladium catalyst and cuprous iodide. The reaction is usually carried out using a base (such as diethylamine, triethylamine or piperidine) in a convenient solvent (such as tetrahydrofuran or dimethylformamide) at a temperature of 20-100°C. Preferably the temperature of the reaction is 25°C. The usual catalyst which is used for the coupling reaction is dichlorobis(triphenylphosphine)-palladium (II).

In an alternative process compounds of formula (I), wherein Z is an acetylene ( $-C\equiv C-$ ) linkage and  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are as defined

above and  $R^1$  is optionally substituted phenyl, can be prepared by treating an acetylene of general formula (IV), (wherein  $R^2, R^3, R^4, R^5, R^6$  and  $R^7$  are as defined above, with a phenyl iodide of general formula ( $R^1I$ ), (wherein  $R^1$  is optionally substituted phenyl), in the presence of a convenient palladium catalyst and cuprous iodide. It is normal to carry out the reaction in the presence of an organic base (such as diethylamine or triethylamine) in a convenient solvent (such as tetrahydrofuran or dimethylformamide) at a temperature of 20-100°C. Preferably the temperature of the reaction is 25°C. The usual palladium catalyst which is used for the coupling reaction is dichlorobis(triphenylphosphine) palladium (II).

The acetylene compound (IV) is prepared by coupling a compound of formula (II) with trimethylsilylacetylene using the already described palladium catalysis and reacting an intermediate of general formula (VI), wherein  $R^2, R^3, R^4, R^5, R^6, R^7$  are as defined above, with potassium hydroxide in methanol as solvent at room temperature.

In a further alternative process compounds of formula (I), wherein Z is an acetylene ( $-C\equiv C-$ ) linkage and  $R^1, R^2, R^3, R^4, R^5, R^6$  and  $R^7$  are as defined above, can be prepared by cross-coupling a bromopyridine of general formula (VII), wherein  $R^1, R^2, R^3, R^4$  are as defined above, with a compound of formula (VIII), wherein  $R^5, R^6$  and  $R^7$  are as defined above and X is bromine or iodine. The coupling reaction can be carried out by treating the bromopyridine (VII) with normal butyl lithium in tetrahydrofuran followed by zinc chloride in diethylether, and then adding tetrakis(triphenylphosphine)palladium (0) with the compound of formula (VIII). The reaction is carried out at temperatures of between -70 and 25°C.

Compounds of formula (I), wherein Z is an olefin ( $-CR^8=CR^{10}-$ ) linkage and  $R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8$  and  $R^{10}$  are as defined above, that is, compounds of general formula (IX) can be prepared by cross-coupling a vinyl halide of general formula (X), wherein  $R^1, R^8, R^{10}$  are as defined above and Hal is a halogen (preferably bromine or iodine), with a halopyridine of general formula (II). The coupling reaction can be carried out by treating the halopyridine (II) with normal butyl lithium in tetrahydrofuran followed by zinc chloride in diethylether and then adding tetrakis(triphenylphosphine)palladium (0) with the vinyl halide (X). The reaction is carried out at temperatures of between -70 and 25°C.

In an alternative process for the production of compounds of general formula (IX), the above cross-coupling reaction can be carried out between a bromopyridine of general formula (XI), wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^8$  and  $R^{10}$  are as defined above, and a compound of formula (VIII).

Compounds of general formula (IX), wherein  $R^8$  and  $R^{10}$  are hydrogen can be prepared by reduction of acetylenes of general formula (XII) wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are as defined above. The reduction can either be achieved by using hydrogen with a suitable catalyst (such as palladium/calcium carbonate/lead oxide (Lindlar catalyst)) or a metal hydride reducing agent (such as lithium aluminium hydride) optionally in the presence of a transition metal catalyst. The reaction is usually carried out at 25°C.

Compounds of formula (I) wherein Z is an ethane  $-CR^8R^9-CR^{10}R^{11}-$  linkage, that is, compounds of general formula (XIII) can be prepared by cross-coupling a halo compound of general formula (XIV), wherein  $R^1$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{11}$  are defined above and Hal is a halogen (preferably bromine or iodine), with a halopyridine of formula (II). The coupling reaction is usually carried out by treating the halo compound (XIV) with normal butyl lithium in tetrahydrofuran followed by zinc chloride in diethyl ether, and then adding tetrakis(triphenylphosphine)palladium (0) with the halopyridine (II).

Alternatively compounds of general formula (XIII) can be prepared by cross-coupling a bromopyridine of general formula (XV), wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{11}$  are defined above, and a compound of formula (VIII).

In an alternative process compounds of general formula (XIII), wherein  $R^9$  and  $R^{11}$  are hydrogen, can be prepared by the reduction of compounds of general formula (IX). The reduction is preferably carried out by hydrogen in the presence of a convenient catalyst, such as palladium on carbon, in a suitable solvent, such as ethanol or ethyl acetate.

The halopyridines (II), bromopyridines (VII), compounds of formula (VIII), bromopyridines (XI) and bromopyridines (XV) are prepared by methods set out in the literature.

The salts of compounds of general formula (I) can be prepared by adding one equivalent of a strong acid such as hydrochloric acid, sulphuric acid or nitric acid, to a solution of (I) in a suitable solvent.

The present invention includes processes for preparing the compounds of formula (I) as herein defined and intermediates of formula (IV) and (VI) as herein before defined.

The compounds are active fungicides and may be used to control one or more of the following pathogens : Pyricularia oryzae on rice; Puccinia recondita, Puccinia striiformis and other rusts on wheat, Puccinia hordei, Puccinia striiformis and other rusts on barley, and rusts on other hosts e.g. coffee, pears, apples, peanuts, vegetables and ornamental plants; Erysiphe graminis (powdery mildew) on barley and wheat and other powdery mildews on various hosts such as Sphaerotheca macularis on hops, Sphaerotheca fuliginea on cucurbits (e.g. cucumber), Podosphaera leucotricha on apple and Uncinula necator on vines; Helminthosporium spp., Rhynchosporium spp., Septoria spp., Pyrenophora spp., Pseudocercospora herpotrichoides and Gaeumannomyces graminis on cereals; Cercospora arachidicola and Cercosporidium personata on peanuts and other Cercospora species on other hosts, for example, sugar beet, bananas, soya beans and rice; Botrytis cinerea (grey mould) on tomatoes, strawberries, vegetables, vines and other hosts; Alternaria spp. on vegetables (e.g. cucumber), oil-seed rape, apples, tomatoes and other hosts; Venturia inaequalis (scab) on apples; Plasmopara viticola on vines; other downy mildews such as Bremia lactucae on lettuce, Peronospora spp. on soybeans, tobacco, onions and other hosts, Pseudoperonospora humuli on hops and Pseudoperonospora cubensis on cucurbits; Phytophthora infestans on potatoes and tomatoes and other Phytophthora spp. on vegetables, strawberries, avocado, pepper, ornamentals, tobacco, cocoa and other hosts; Thanatephorus cucumeris on rice and other Rhizoctonia species on various hosts such as wheat and barley, vegetables, cotton and turf.

Some of the compounds show a broad range of activities against fungi in vitro. They may also have activity against various post-harvest diseases of fruit (e.g. Penicillium digitatum and italicum and Trichoderma viride on oranges, Gloeosporium musarum on bananas and Botrytis cinerea on grapes).

Further, some of the compounds may be active as seed dressings against pathogens including Fusarium spp., Septoria spp., Tilletia spp., (bunt, a seed-borne disease of wheat), Ustilago spp. and Helminthosporium spp. on cereals, Rhizoctonia solani on cotton and Pyricularia oryzae on rice.

The compounds may move acropetally/locally in plant tissue. Moreover, the compounds may be volatile enough to be active in the vapour phase against fungi on the plant.

The invention therefore provides a method of combating fungi which comprises applying to a plant, to a seed of a plant or to the locus of the

plant or seed a fungicidally effective amount of a compound as hereinbefore defined, or a composition containing the same.

The compounds may be used directly for agricultural purposes but are more conveniently formulated into compositions using a carrier or diluent. The invention thus provides fungicidal compositions comprising a compound as hereinbefore defined and an acceptable carrier or diluent therefor. It is preferred that these compositions comprise 0.0001 to 95%, more preferably 0.001 to 60%, even more preferably 0.01 to 1%, of the compound.

The compounds can be applied in a number of ways. For example, they can be applied, formulated or unformulated, directly to the foliage of a plant, to seeds or to other medium in which plants are growing or are to be planted, or they can be sprayed on, dusted on or applied as a cream or paste formulation, or they can be applied as a vapour or as slow release granules.

Application can be to any part of the plant including the foliage, stems, branches or roots, or to soil surrounding the roots, or to the seed before it is planted, or to the soil generally, to paddy water or to hydroponic culture systems. The invention compounds may also be injected into plants or sprayed onto vegetation using electrodynamic spraying techniques or other low volume methods.

The term "plant" as used herein includes seedlings, bushes and trees. Furthermore, the fungicidal method of the invention includes preventative, protectant, prophylactic and eradicant treatments.

The compounds are preferably used for agricultural and horticultural purposes in the form of a composition. The type of composition used in any instance will depend upon the particular purpose envisaged.

The compositions may be in the form of dustable powders or granules comprising the active ingredient (invention compound) and a solid diluent or carrier, for example, fillers such as kaolin, bentonite, kieselguhr, dolomite, calcium carbonate, talc, powdered magnesia, fuller's earth, gypsum, diatomaceous earth and china clay. Such granules can be preformed granules suitable for application to the soil without further treatment. These granules can be made either by impregnating pellets of filler with the active ingredient or by pelleting a mixture of the active ingredient and powdered filler. Compositions for dressing seed may include an agent (for example, a mineral oil) for assisting the adhesion of the composition to the seed; alternatively the active ingredient can be formulated for seed dressing purposes using an organic solvent (for example, N-methylpyrrol-

idone, propylene glycol or N,N-dimethylformamide). The compositions may also be in the form of wettable powders or water dispersible granules comprising wetting or dispersing agents to facilitate the dispersion in liquids. The powders and granules may also contain fillers and suspending agents.

Emulsifiable concentrates or emulsions may be prepared by dissolving the active ingredient in an organic solvent optionally containing a wetting or emulsifying agent and then adding the mixture to water which may also contain a wetting or emulsifying agent. Suitable organic solvents are aromatic solvents such as alkylbenzenes and alkyl naphthalenes, ketones such as cyclohexanone and methylcyclohexanone, chlorinated hydrocarbons such as chlorobenzene and trichlorethane, and alcohols such as benzyl alcohol, furfuryl alcohol, butanol and glycol ethers.

Suspension concentrates of largely insoluble solids may be prepared by ball or bead milling with a dispersing agent with a suspending agent included to stop the solid settling.

Compositions to be used as sprays may be in the form of aerosols wherein the formulation is held in a container under pressure of a propellant, e.g. fluorotrichloromethane or dichlorodifluoromethane.

The invention compounds can be mixed in the dry state with a pyrotechnic mixture to form a composition suitable for generating in enclosed spaces a smoke containing the compounds.

Alternatively, the compounds may be used in micro-encapsulated form. They may also be formulated in biodegradable polymeric formulations to obtain a slow, controlled release of the active substance.

By including suitable additives, for example additives for improving the distribution, adhesive power and resistance to rain on treated surfaces, the different compositions can be better adapted for various utilities.

The invention compounds can be used as mixtures with fertilisers (e.g. nitrogen-, potassium- or phosphorus-containing fertilisers). Compositions comprising only granules of fertiliser incorporating, for example coated with, the compound are preferred. Such granules suitably contain up to 25% by weight of the compound. The invention therefore also provides a fertiliser composition comprising a fertiliser and the compound of general formula (I) or a salt or metal complex thereof.

Wettable powders, emulsifiable concentrates and suspension concentrates will normally contain surfactants, e.g. a wetting agent,

dispersing agent, emulsifying agent or suspending agent. These agents can be cationic, anionic or non-ionic agents.

Suitable cationic agents are quaternary ammonium compounds, for example, cetyltrimethylammonium bromide. Suitable anionic agents are soaps, salts of aliphatic monoesters of sulphuric acid (for example, sodium lauryl sulphate), and salts of sulphonated aromatic compounds (for example, sodium dodecylbenzenesulphonate, sodium, calcium or ammonium lignosulphonate, butylnaphthalene sulphonate, and a mixture of sodium diisopropyl- and triisopropylnaphthalene sulphonates).

Suitable non-ionic agents are the condensation products of ethylene oxide with fatty alcohols such as oleyl or cetyl alcohol, or with alkyl phenols such as octyl- or nonylphenol and octylcresol. Other non-ionic agents are the partial esters derived from long chain fatty acids and hexitol anhydrides, the condensation products of the said partial esters with ethylene oxide, and the lecithins. Suitable suspending agents are hydrophilic colloids (for example, polyvinylpyrrolidone and sodium carboxymethylcellulose), and swelling clays such as bentonite or attapulgite.

Compositions for use as aqueous dispersions or emulsions are generally supplied in the form of a concentrate containing a high proportion of the active ingredient, the concentrate being diluted with water before use. These concentrates should preferably be able to withstand storage for prolonged periods and after such storage be capable of dilution with water in order to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. The concentrates may conveniently contain up to 95%, suitably 10-85%, for example 25-60%, by weight of the active ingredient. After dilution to form aqueous preparations, such preparations may contain varying amounts of the active ingredient depending upon the intended purpose, but an aqueous preparation containing 0.0005% or 0.01% to 10% by weight of active ingredient may be used.

The compositions of this invention may contain other compounds having biological activity, e.g. compounds having similar or complementary fungicidal activity or which possess plant growth regulating, herbicidal or insecticidal activity.

A fungicidal compound which may be present in the composition of the invention may be one which is capable of combating ear diseases of cereals (e.g. wheat) such as Septoria, Gibberella and Helminthosporium spp., seed

and soil-borne diseases and downy and powdery mildews on grapes and powdery mildew and scab on apple, etc. By including another fungicide, the composition can have a broader spectrum of activity than the compound of general formula (I) alone. Further the other fungicide can have a synergistic effect on the fungicidal activity of the compound of general formula (I). Examples of fungicidal compounds which may be included in the composition of the invention are (+)-cis-1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol, (2RS,3RS)-1-[3-(2-chlorophenyl)-2-(4-fluorophenyl)oxiran-2-ylmethyl]-1H-1,2,4-triazole, (RS)-1-aminopropylphosphonic acid, (RS)-4-(4-chlorophenyl)-2-phenyl-2-(1H-1,2,4-triazol-1-ylmethyl)butyronitrile, (Z)-N-but-2-enyloxymethyl-2-chloro-2',6'-diethylacetanilide, 1-(2-cyano-2-methoxyiminoacetyl)-3-ethyl urea, 3-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-yl)quinazolin-4(3H)-one, 4-(2,2-difluoro-1,3-benzodioxol-4-yl)pyrrole-3-carbonitrile, 4-bromo-2-cyano-N,N-dimethyl-6-trifluoromethylbenzimidazole-1-sulphonamide, 5-ethyl-5,8-dihydro-8-oxo(1,3)-dioxol-(4,5-g)quinoline-7-carboxylic acid,  $\alpha$ -[N-(3-chloro-2,6-xylyl)-2-methoxyacetamido]- $\gamma$ -butyrolactone, aldimorph, anilazine, benalaxyl, benomyl, bixoxazol, binapacryl, bitertanol, blasticidin S, bromuconazole, bupirimate, buthiobate, captafol, captan, carbendazim, carboxin, chlorbenzthiazole, chloroneb, chlorothalonil, chlorozolinate, copper containing compounds such as copper oxychloride, copper sulphate and Bordeaux mixture, cycloheximide, cymoxanil, cyproconazole, cyprofuram, di-2-pyridyl disulphide 1,1'-dioxide, dichlofluanid, dichlone, diclobutrazol, diclomezine, dicloran, difenoconazole, dimethamorph, dimethirimol, diniconazole, dinocap, ditalimfos, dithianon, dodemorph, dodine, edifenphos, etaconazole, ethirimol, ethyl (Z)-N-benzyl-N-([methyl(methyl-thioethylideneamino-oxycarbonyl)amino]thio)- $\beta$ -alaninate, etridiazole, fenapanil, fenarimol, fenfuram, fencpiclonil, fenpropidin, fenpropimorph, fentin acetate, fentin hydroxide, flutolanil, flutriafol, flusilazole, folpet, fosetyl-aluminium, fuberidazole, furalaxyl, furconazole-cis, guazatine, hexaconazole, hydroxyisoxazole, imazalil, imibenconazole, iprobenfos, iprodione, isoprothiolane, kasugamycin, mancozeb, maneb, mepanipyrim, mepronil, metalaxyl, methfuroxam, metsulfosax, myclobutanil, neoasozin, nickel dimethyldithiocarbamate, nitrothal-isopropyl, nuarimol, ofurace, organomercury compounds, oxadixyl, oxycarboxin, pefurazoate, penconazole, pencycuron, phenazin oxide, phthalide, polyoxin D, polyram, probenazole, prochloraz, procymidone, propamocarb, propiconazole, propineb, prothiocarb, pyrazophos, pyrifenoxy, pyroquilon, pyroxyfur, pyrrolnitrin, quinomethionate, quintozone, streptomycin, sulphur,

tebuconazole, techlofthalam, tecnazene, tetraconazole, thiabendazole, thicyofen, thiophanate-methyl, thiram, tolclofos-methyl, triacetate salt of 1,1'-iminodi(octamethylene)diguanidine, triadimefon, triadimenol, triaz-butyl, tricyclazole, tridemorph, triforine, validamycin A, vinclozolin, zarilamid and zineb. The compounds of general formula (I) can be mixed with soil, peat or other rooting media for the protection of plants against seed-borne, soil-borne or foliar fungal diseases.

Suitable insecticides which may be incorporated in the composition of the invention include buprofezin, carbaryl, carbofuran, carbosulfan, chlorpyrifos, cycloprothrin, demeton-s-methyl, diazinon, dimethoate, ethofenprox, fenitrothion, fenobucarb, fenthion, formothion, isoprocab, isoxathion, monocrotophos, phenthoate, pirimicarb, propaphos and XMC.

Plant growth regulating compounds are compounds which control weeds or seedhead, formation, or selectively control the growth of less desirable plants (e.g. grasses).

Examples of suitable plant growth regulating compounds for use with the invention compounds are 3,6-dichloropicolinic acid, 1-(4-chlorophenyl)-4,6-di-methyl-2-oxo-1,2-dihydropyridine-3-carboxylic acid, methyl-3,6-dichloroanisate, abscisic acid, asulam, benzoylprop-ethyl, carbetamide, daminozide, difenzoquat, dikegulac, ethephon, fenpentezol, fluoridamid, glyphosate, glyphosine, hydroxybenzonitriles (e.g. bromoxynil), inabenfide, isopyrimol, long chain fatty alcohols and acids, maleic hydrazide, mefluidide, morphactins (e.g. chlorfluoroecol), paclobutrazol, phenoxyacetic acids (e.g. 2,4-D or MCPA), substituted benzoic acid (e.g. triiodobenzoic acid), substituted quaternary ammonium and phosphonium compounds (e.g. chloromequat, chlorphonium or mepiquatchloride), tecnazene, the auxins (e.g. indoleacetic acid, indolebutyric acid, naphthylacetic acid or naphthoxyacetic acid), the cytokinins (e.g. benzimidazole, benzyladenine, benzylaminopurine, diphenylurea or kinetin), the gibberellins (e.g. GA<sub>3</sub>, GA<sub>4</sub> or GA<sub>7</sub>) and triapenthenol.

The following Examples illustrate the invention. Throughout the examples reactions involving water-sensitive intermediates were performed under atmospheres of nitrogen. Unless otherwise stated, chromatography was performed using silica gel as the stationary phase. DISPERSOL and TWEEN, where used in the Examples, are Trade or Service Marks.

EXAMPLE 1

This Example illustrates the preparation of 2-(6-phenylethynyl-2-pyridinyl)-4-methylpyrimidine (Compound No. 1 in Table I).

Stage 1

A solution of normal butyl lithium (28.6ml, 0.0714 moles of a 2.5M solution in hexane) was added dropwise to a solution of 2,6-dibromopyridine (15.8g, 0.067 moles) in dry tetrahydrofuran (100ml) at -78°C under a nitrogen atmosphere. The mixture was stirred for 0.5 hours at -78°C after the addition was complete and a solution of zinc chloride (205ml, 0.2047 moles of a 1.0M solution in diethyl ether) added dropwise so that the temperature was kept below -60°C. A white precipitate formed and the solution was stirred at -78°C for a further 0.5 hours.

Tetrakis(triphenylphosphine)palladium (0) (1.1g, 0.00095 moles) was added, followed by 2-bromo-4-methyl-pyrimidine (8.23g, 0.0476 moles) in dry tetrahydrofuran (100ml). After complete addition the reaction mixture was refluxed for 4 hours, cooled to room temperature, poured into a 10% solution of ethylenediamine-tetraacetic acid and extracted with methylene chloride. The combined methylene chloride extracts were washed, dried and the solvent removed to give a dark oil which was purified by column chromatography (silica gel eluted with *t*-butyl methyl ether) giving 2-(6-bromo-2-pyridinyl)-4-methylpyrimidine (3.6, 30%) as a yellow solid. This was used in the next stage without further recrystallisation.

Stage 2

Copper (I) iodide (0.057g, 0.0003 moles) was added portionwise to a stirred suspension of 2-(6-bromo-2-pyridinyl)-4-methylpyrimidine (2.5g, 0.01 moles), phenylacetylene (2.04g, 0.02 moles) and dichlorobis-(triphenylphosphine)palladium (II) (0.105g, 0.00015 moles) in triethylamine (90ml) and dry tetrahydrofuran (60ml) under a nitrogen atmosphere at 20°C. The reaction mixture was stirred at room temperature for 20 hours and the solvent removed to give a dark residue. This residue was dissolved in ethyl acetate, washed and dried over anhydrous magnesium sulphate. Removal of the solvent gave a black oil which was purified by column chromatography (silica gel eluted with dichloromethane : tetrahydrofuran 98:2) to give the title compound (1.47g, 54%) as an orange solid mp 83-85°C.

EXAMPLE 2

This Example illustrates the preparation of 2-(6-styryl-2-pyridinyl)-4-methylpyrimidine (Compound No. 2 in Table I).

Lithium aluminium hydride (0.8ml of 1.0M solution in tetrahydrofuran) was added dropwise to a mixture of 2-(6-phenylethynyl-2-pyridinyl)-4-methylpyrimidine (0.4g, 0.0015 moles) and ferrous chloride (0.15g, 0.00075 moles) in dry tetrahydrofuran (1.0ml) at  $-78^{\circ}\text{C}$  under a nitrogen atmosphere. After complete addition, the solution was stirred at  $-78^{\circ}\text{C}$  for 0.5 hours and allowed to warm up to room temperature. The reaction mixture was poured carefully into water, extracted with ethyl acetate, washed and dried over anhydrous magnesium sulphate. Removal of the solvent gave a brown oil which was purified by column chromatography (silica gel eluted with diethyl ether) to give the title compound (0.054g, 13%) as a yellow oil which is a mixture of isomers (trans:cis ratio of 1:2).

The following are examples of compositions suitable for agricultural and horticultural purposes which can be formulated from the compounds of the invention. Such compositions form another aspect of the invention. Percentages are by weight.

#### EXAMPLE 3

This Example illustrates the preparation of 2-(6-(4-chlorophenylethynyl)pyrid-2-yl)-4-methylpyrimidine (Compound No. 3 in Table I).

##### Step 1

Dichlorobis(triphenylphosphine)palladium (II) (0.66g, 0.00094 moles) and copper (I) iodide (0.36g, 0.0019 moles) were added to a solution of chloro-4-iodobenzene (6.0g, 0.0252 moles), trimethylsilylacetylene (4.9g, 0.0504 moles), and triethylamine (5.2cm<sup>3</sup>, 3.8g, 0.038 moles) in N,N-dimethylformamide (60cm<sup>3</sup>), under an atmosphere of nitrogen. The reaction was stirred, under nitrogen, at room temperature for 40 minutes, giving a dark brown mixture.

This was poured onto water (200cm<sup>3</sup>), a 10% solution of ethylenediamine tetraacetic acid sodium salt (100cm<sup>3</sup>) was added and the mixture was extracted with ethyl acetate (100cm<sup>3</sup>). The organic extract was dried and the solvent removed to give a dark oil which was purified by preparative high performance liquid chromatography, (silica gel eluted with hexane) giving chloro-4-trimethylsilylethynylbenzene as a yellow solid. This was used in the next step without further recrystallisation. Yield 3.0g (58%).

<sup>1</sup>H NMR: 7.0 (2H,d), 7.25 (2H,d), 0.0 (not integrated but should be 9H,s)ppm.

##### Step 2

Anhydrous potassium carbonate (0.9g, 0.0065 moles) was added to a solution of chloro-4-trimethylsilylethynylbenzene (3.0g, 0.014 moles) in

methanol (60cm<sup>3</sup>) under an atmosphere of nitrogen. The reaction was stirred, under nitrogen, at room temperature for 4 hours. The solvent was removed, the solid residue was triturated with ether and then dichloromethane and filtered. The organic filtrates were combined and evaporated to dryness giving chloro-4-ethynylbenzene as a brown solid. This was used in the next step without any further purification yield 1.1g (58%). <sup>1</sup>H NMR δ7.3 (2H, d), 7.4 (2H, d), 3.1 (1H, s)ppm.

### Step 3

A solution of chloro-4-ethynylbenzene (0.48g, 0.0035 moles) in dry tetrahydrofuran (8cm<sup>3</sup>) was added to a solution of 2-(6-bromo-2-pyridinyl)-4-methyl pyrimidine (0.44g, 0.00176 moles) in dry tetrahydrofuran (8cm<sup>3</sup>) under a nitrogen atmosphere. Dry triethylamine (12cm<sup>3</sup>), dichlorobis(triphenylphosphine) palladium (II) (0.02g, 0.000028 moles) and copper (I) iodide (99.999%) (0.02g, 0.0001 moles) were then added and the mixture was stirred at room temperature, under nitrogen, for 22 hours.

The solvent was removed and the residue was partitioned between ethyl acetate and water. The organic phase was dried and evaporated to give a brown solid residue which was purified by preparative thin layer chromatography using either as eluent. The title compound was obtained as as a pale brown solid. Yield 0.25g (46%).

### EXAMPLE 4

This Example illustrates the preparation of 2-[6-(3,3-dimethylbut-1-ynyl)-2-pyridinyl]-4-methylpyrimidine (Compound No. 99 in Table I).

2-(6-Bromo-2-pyridinyl)-4-methylpyrimidine (0.5g, 0.002 moles) [prepared as in Stage 1 of Example 1] was dissolved in dry tetrahydrofuran and 3,3-dimethylbut-1-yne (0.33g, 0.004 moles) was added, followed by dry triethylamine (12cm<sup>3</sup>), dichlorobis(triphenylphosphine)palladium (II) (0.02g, 0.000028 moles). The reaction was left at room temperature for 17 hours then heated at 40° for 4 hours. More 3,3-trimethylbut-1-yne (0.33g, 0.5cm<sup>3</sup>, 0.004 moles) was added, after which the reaction mixture was heated at reflux for 2 hours and left to stand at room temperature overnight. More 3,3-dimethylbut-1-yne (0.67g, 0.008 moles), palladium (II) catalyst (0.02g, 0.000028 moles) and copper (I) iodide (0.01g, 0.000053 moles) were added and reaction was heated at reflux for 3 hours. Piperidine (5cm<sup>3</sup>) was added, giving a dark brown solution which was heated at reflux for 1½ hours.

The mixture was evaporated to dryness under reduced pressure and the residue was partitioned between ethyl acetate and water. The organic phase

was dried, then evaporated to give a dark brown viscous oil, which was purified by preparative thin layer chromatography on silica gel with ether as eluent. The title compound was obtained as a pale brown oil which solidified on standing (0.12g, 24%, mpt 98-100°C).

#### EXAMPLE 5

An emulsifiable concentrate is made up by mixing and stirring the ingredients until all are dissolved.

Compound No. 1 of Table I		10%
Benzyl alcohol	30%	
Calcium dodecylbenzenesulphonate	5%	
Nonylphenolethoxylate (13 moles ethylene oxide)	10%	
Alkyl benzenes	45%	

#### EXAMPLE 6

The active ingredient is dissolved in methylene dichloride and the resultant liquid sprayed on to the granules of attapulgite clay. The solvent is then allowed to evaporate to produce a granular composition.

Compound No. 2 of Table I	5%
Attapulgite granules	95%

#### EXAMPLE 7

A composition suitable for use as a seed dressing is prepared by grinding and mixing the three ingredients.

Compound No. 1 of Table I	50%
Mineral oil	2%
China clay	48%

#### EXAMPLE 8

A dustable powder is prepared by grinding and mixing the active ingredient with talc.

Compound No. 1 of Table I	5%
Talc	95%

#### EXAMPLE 9

A suspension concentrate is prepared by ball milling the ingredients to form an aqueous suspension of the ground mixture with water.

Compound No. 1 of Table I	40%
Sodium lignosulphonate	10%
Bentonite Clay	1%
Water	49%

This formulation can be used as a spray by diluting into water or

applied directly to seed.

#### EXAMPLE 10

A wettable powder formulation is made by mixing together and grinding the ingredients until all are thoroughly mixed.

Compound No. 1 of Table I	25%
Sodium lauryl sulphate	2%
Sodium lignosulphonate	5%
Silica	25%
China clay	43%

#### EXAMPLE 11

The compounds were tested against a variety of foliar fungal diseases of plants. The technique employed was as follows.

The plants were grown in John Innes Potting Compost (No 1 or 2) in 4 cm diameter minipots. The test compounds were formulated by bead milling with aqueous DISPERSOL T which was diluted to the required concentration immediately before use. For the foliage diseases, the formulations (100 ppm active ingredient) were sprayed onto the foliage and applied to the roots of the plants in the soil. The sprays were applied to maximum retention and the root drenches to a final concentration equivalent to approximately 40 ppm a.i. in dry soil. TWEEN 20, to give a final concentration of 0.05%, was added when the sprays were applied to cereals.

For most of the tests the compound was applied to the soil (roots) and to the foliage (by spraying) one or two days before the plant was inoculated with the disease. An exception was the test on Erysiphe graminis in which the plants were inoculated 24 hours before treatment. Foliar pathogens were applied by spray as spore suspensions onto the leaves of test plants. After inoculation, the plants were put into an appropriate environment to allow infection to proceed and then incubated until the disease was ready for assessment. The period between inoculation and assessment varied from four to fourteen days according to the disease and environment.

The disease control was recorded by the following grading:




- 4 = no disease
- 3 = trace -5% of disease on untreated plants
- 2 = 6-25% of disease on untreated plants
- 1 = 26-59% of disease on untreated plants
- 0 = 60-100% of disease on untreated plants





The results are shown in Table III.




TABLE III



Compound	Puccinia recondita (Wheat)	Erysiphe graminis tritici (Wheat)	Venturia inaequalis (Apple)	Pyricularia oryzae (Rice)	Thanetophorus cucumeris (Rice)	Septoria nodorum	Plasmopara viticola (Vine)
1	3	4	4	4	-	0	1
2	0*	0*	4*	3*	-	3*	0*
3	-	2	2	4	0	0	3
6	-	0	4	4	0	0	1
10	0	0	0	1	0	0	2
95	-	0	1	2	0	0	4
96	-	0	3	3	0	3	4
97	0	0	4	3	1	0	3

\* data generated at 25 ppm.

CHEMICAL FORMULAE  
(in description)









CLAIMS

1. A compound of general formula (I):




wherein  $R^1$  is an optionally substituted phenyl group,  $C_{3-6}$  cycloalkyl optionally substituted by alkyl or alkyl;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  alkoxy or  $NR'R''$ ; Z is a linking group and is  $-C\equiv C-$ ,  $-CR^8=CR^{10}-$  or  $-CR^8R^9-CR^{10}R^{11}-$ ;  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{11}$ ,  $R'$  and  $R''$  are independently hydrogen or  $C_{1-4}$  alkyl; provided that when  $R^1$  is alkyl Z is not  $-CR^8R^9-CR^{10}R^{11}-$ ; and acid addition salts or metal complexes thereof.

2. A compound of general formula (I):




wherein  $R^1$  is an optionally substituted phenyl group;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  haloalkyl or  $C_{1-4}$  alkoxy; Z is a linking group and is  $-C\equiv C-$ ,  $-CR^8=CR^{10}-$  or  $-CR^8R^9-CR^{10}R^{11}-$ ;  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  are independently hydrogen or  $C_{1-4}$  alkyl; and acid addition salts or metal complexes thereof.

3. A compound as claimed in claim 1 or 2 having the general formula (I):



wherein  $R^1$  is an optionally substituted phenyl group,  $C_{3-6}$  cycloalkyl optionally substituted by alkyl or alkyl;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  alkoxy or  $NR'R''$ ; Z is  $-CR^8=CR^{10}-$  or  $-C\equiv C-$ ;  $R^8$ ,  $R^{10}$ ,  $R'$  and  $R''$  are independently hydrogen or  $C_{1-4}$  alkyl; and acid addition salts or metal complexes thereof.


4. A compound as claimed in claim 1, 2 or 3 wherein  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$  and  $R^7$  are all hydrogen and  $R^5$  is  $C_{1-4}$  alkyl.
5. A compound as claimed in claim 1, 2, 3 or 4 having the general formula (I):



wherein  $R^1$  is phenyl optionally substituted with halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkoxy,  $C_{2-4}$  alkenyl,  $C_{2-4}$  alkynyl,  $C_{1-4}$  haloalkyl or phenyl; Z is  $-C\equiv C-$ ;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$  and  $R^7$  are all


hydrogen;  $R^5$  is  $C_{1-4}$  alkyl; and acid addition salts or metal complexes thereof.

6. A compound as claimed in claim 1 or 2 having the general formula (I):




wherein  $R^1$  is phenyl optionally substituted with halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  haloalkoxy or phenyl, or  $R^1$  is cyclohexyl or tert-butyl; Z  $-CH=CH-$  or  $-C\equiv C-$ ;  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^6$  and  $R^7$  are all hydrogen; and  $R^5$  is  $C_{1-4}$  alkyl.

7. A process for the preparation of a compound of general formula (I) as defined in claim 1 comprising:
- a) cross-coupling a compound of formula (II):




wherein Hal is a halogen atom and  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are as defined in claim 1, with a compound of formula (III), (X) or (XIV):




wherein  $R^1$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  are as defined in claim 1 and Hal as a halogen;

or

b) cross-coupling a compound of formula (XVI):




wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and Z are as defined in claim 1; with a bromopyrimidine of general formula (VIII):



wherein  $R^5$ ,  $R^6$  and  $R^7$  are as defined in claim 1;


or

c) to prepare a compound of formula (I) wherein Z is  $-C\equiv C-$ , treating a compound of formula (IV):



wherein  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are as defined in claim 1, with a phenyl iodide of general formula  $R^1I$ , wherein  $R^1$  is as defined in claim 1.

8. The intermediate compound of general formula (IV):



wherein  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are as defined in claim 1.

9. A fungicidal composition comprising a fungicidally effective amount of a compound according to claim 1 and a fungicidally acceptable carrier or diluent therefor.
10. A method of combating fungi which comprises applying to plants, to the seeds of plants or to the locus of the plants or seeds, a compound according to claim 1 or a composition according to claim 9.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 91/01853

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C07D401/04; A01N43/54		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.Cl. 5	C07D	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
X	EP,A,0 234 104 (SUMITOMO CHEMICAL COMPANY LIMITED) 2 September 1987 see claims; examples	1,9,10
P,X	EP,A,0 407 899 (HOECHST AKTIENGESELLSCHAFT) 16 January 1991 see claims	1,9,10
P,X	WO,A,9 107 399 (HOECHST AKTIENGESELLSCHAFT) 30 May 1991 see claims	1,9,10
<p><sup>10</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
09 JANUARY 1992	30.01.92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	DE JONG B.S.	

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO. GB 9101853  
SA 52637**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 09/01/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0234104	02-09-87	JP-A- 62169778	25-07-87
		US-A- 4752608	21-06-88
		JP-A- 63099068	30-04-88
EP-A-0407899	16-01-91	DE-A- 3922735	24-01-91
WO-A-9107399	30-05-91	DE-A- 3937285	16-05-91
		AU-A- 6886891	13-06-91
		CN-A- 1051558	22-05-91